
CITES APPLICATION FOR REGISTRATION AND CCREDITATION OF OPERATION BREEDING APPENDIX I SPECIES Yellow-Naped Amazon (Amazona ochrocephala auropalliata) FOR COMMERCIAL PURPOSES (Res. Conf. 12.10)



Birds International, Inc. No. 1 Sto Nino St. Alemars Subd. Commonwealth Heights, Fairview, Quezon City, Philippines

INTRODUCTION

This proposal is being submitted to the CITES Secretariat-Switzerland through the CITES Management Authority of the Philippines, the Protected Areas and Wildlife Bureau (PAWB), for the purpose of gaining accreditation for the possible trade of Birds International, Inc.'s captive-bred, second and succeeding generations of Yellow-naped Amazon (Amazona ochrocephala auropalliata).

Birds International, Inc. (BII) is engaged in the conservation and propagation of the world's rare and exotic species of psittacines, 95% of which are non-Philippine birds. It operates its activities on a six-hectare farm in Quezon City, Philippines. BII is a CITES registered breeding facility (A-PH-501) breeding Appendix 1 species of Golden Conure (Aratinga guarouba) for commercial purposes.

While most breeding and conservation efforts done are government-initiated and sponsored, BII was founded privately by Antonio M. de Dios. What started as a pure hobby became a well-organized, systematic and highly technical operation to breed and conserve, which ultimately culminated in the establishment of BII.

Upon its establishment in the early 1970s, BII's principal aim was to undertake biological and conservation studies for future breeding efforts. Also, a continuous exchange of information was established with other international aviculturists.

An influx of investments followed to be able to achieve the desired breeding conditions and provide the expertise required for the successful propagation of the birds. Necessary facilities such as a veterinary hospital with twenty rooms to accommodate various cases; a laboratory with high-powered microscopes, a serum analyzer unit, a mechanical convection unit; breeding and non-breeding cages and aviaries of various designs and sizes; a 45 air-conditioned room nursery; a fully insulated hatchery with more than 40 units of Grumbach egg incubators and hatchers; dietary centers; water filtration systems, sprinkler systems for birds' artificial shower and staff quarters were constructed. In addition, hundreds of animal incubators and other needed equipment, feeds and medicines were imported.

BII is jointly managed by Mr. Antonio de Dios and his daughter, Regina. They have a staff of one hundred thirty (130) people consisting of college graduates with degrees in veterinary medicine, medical technology, animal husbandry, agriculture and zoology; and a support group of maintenance people consisting of gardeners, janitors, carpenters, masons, electricians and welders. Functional groups carry out specialized activities such as hospital management, nursery management, management of facilities and equipment and the likes.

In its more than thirty years of bird conservation and propagation, BII has made substantial contributions to the accumulation of technical knowledge on bird farming. It has achieved breeding success in its Appendix I and II species, most notably the Spix's Macaw <u>Cyanopsitta spixili</u>

The Philippines became an active member of the Convention on International Trade of Endangered Species Flora and Fauna (CITES) in 1981. With its strong adherence to the CITES regulation of restricting the trade of wildlife species only to those that are born and bred in captivity, BII was registered as a preferred pioneer industry with the Board of Investments of the Philippines (BOI) Certificate of Registration No. 83-598 on November 16, 1983. On August 14, 1985, BII was granted Wildlife Permit No. 3 by the then Bureau of Forest Development, which allowed Birds International, Inc to export its captive-bred birds.

Net revenue generated by BII is flowed back to its operation. We continually maintain and upgrade our facilities, we hold training and seminars for our staff conducted by international professionals and we also sponsor and undertake training programs for both local and foreign veterinarians from various zoos like the Sao Paolo Zoo in Brazil, Criadouro Chaparral also in Brazil and the Philippine Eagle Conservation Program. We have allowed members of some Asian parks like the Jurong Bird Park in Singapore to observe our system of operation. We engage in exchanging ideas and information exchange with zoos and aviculturists from around the world to give the best care possible to these beautiful avian species. In the end, if such a need should arise, we are willing to be a part of their reintroduction to the wild.

 NAME AND ADDRESS OF THE OWNER AND THE MANAGER OF THE CAPTIVE BREEDING OPERATION.

Birds International Incorporated (BII)
No. 1 Sto Nino St. Alemars Subd.
Commonwealth Heights, Fairview, Quezon City Philippines
Website Address: www.birdsinternational.net
e-mail: info@birdsinternational.net

Mr. Antonio M. de Dios – Founder-President Ms. Ma. Regina de Dios-Jardinel - Manager

- 2. DATE OF ESTABLISHMENT: February 1975
- 3. SPECIES BRED: Yellow-Naped Amazon (Amazona ochrocephala auropalliata)¹
- DETAILS OF THE NUMBER AND AGE (IF KNOWN OR APPROPRIATE) OF MALES AND FEMALES THAT COMPRISE THE PARENTAL BREEDING STOCKS.
 (Evidence of legal acquisition of each male and female including receipts, CITES documents, capture permits, etc)

The company (BII, Philippines²) acquired a total of forty heads (40) Yellownaped Amazon³ with unknown age from Singapore⁴. From this total, twenty four (24) heads or twelve (12.12) pairs became the parental breeding pairs, while sixteen heads (16) died during the quarantine period and. Refer to Annex A for the details.

 OPERATIONS LOCATED WITHIN RANGE STATES MUST PRODUCE EVIDENCE THAT THE PARENTAL STOCK WAS OBTAINED IN ACCORDANCE WITH THE RELEVANT NATIONAL LAWS (e.g capture permits, receipts, etc) or, if imported, in accordance with the provisions of the Convention (e.g. receipts, CITES documents, etc)

Not Applicable Birds International Inc. (BII) is located in a non-range state.

6. OPERATIONS LOCATED IN NON-RANGE STATES MUST PRODUCE EVIDENCE THAT THE ANIMALS COMPRISING THE PARENTAL STOCK ARE: a) are preconvention specimen (relevant dated receipts or other acceptable proof of lawful acquisition) b) have been derived from pre-convention specimens (relevant dated receipts or other acceptable proof of lawful acquisition) c) or were acquired from the range State in accordance with the provisions of the Conventions.

All Yellow-Naped Amazon specimens imported by BII from Singapore (non-range State) in 1980 were allowed and covered by import permits issued by BAI (Bureau of Animal Industry). From the total acquisition of forty (40) specimens, only twelve (12.12) pairs was set-up to become the parental breeding stocks. Refer to Annex B for the details. Refer to Exhibit I for the BAI issued Import Permits.

Acceded to CITES on February 28, 1987

Listed to Appendix II on June 06, 1981 Elevated to Appendix I on February 13, 2003

Elevated to Appendix I on February 13, 2003 Acceded to CITES on November 16, 1981

Acceded to CITES on Movember 10, 1901 Acquired through BAI (Bureau of Animal Industry) Import Permit No.P-153 dated 1980-06-26

 CURRENT STOCKS (numbers, by sex and age, of progeny held in addition to the parental breeding stock above)

The company's current stock of thirty-nine (39) specimens are composed of F1 breeding pairs at sixteen pairs (16.16) and seven (7) F2 progenies. Please refer to table below.

Total Current Stock of Yellow-Naped Amazon

Age		Total		
(Years)	Male	Female	Unsexed	
		0	0	0
1	0	0	0	0
4	2	2	0	4
6	0	1	0	1
8	3	3	0	6
10	7	8	0	15
12	3	4	0	7
14	3	2	0	5
16	0	1	0	1
18	0	0	0	0
20	0	0	0	0
Total	18	21	0	39

8. INFORMATION ON THE PERCENTAGE OF MORTALITIES IN THE DIFFERENT AGE GROUPS AND WHERE POSSIBLE, BETWEEN MALES AND FEMALES.


From the total production of forty-five(45) F1-generation progenies, twenty-nine percent (29%) died, while seventy-one percent (71%) are the remaining live progenies. Refer to Annex C the details

From the total of one hundred eight (108) F2-generation progenies, one percent (1%) died, while ninety-nine percent (99%) are the remaining live progenies. Refer to Annex C details.


 DOCUMENTATION SHOWING THAT THE SPECIES HAS BEEN BRED TO SECOND GENERATION BREEDING OFFSPRING (F2) AT THE FACILITY AND THE DESCRIPTION OF METHOD USED.

When Birds International started Yellow-naped Amazon, the company decided to practice the CLOSED CYCLE BREEDING AND PRODUCTION SYSTEM. This production system requires an initial take from the wild but after the initial establishment no wild specimen will be added or taken from the wild to augment the existing parental breeding pairs. Likewise, the company follows a SEGREGATION STRATEGY of progenies coming from various generations. Segregation will ensure grouping of progenies coming from the same generation and will eliminate chances of mixing individuals from the different generations therefore making pairing of similar generation easier and the realization of F2 and F3 generation attainable.

Diagram below will show how BII was able to achieve F1 and F2 generation from two (2) pairs of Parental Breeding Pair.



Presented below is the CLOSED CYCLE BREEDING AND PRODUCTION SYSYTEM for the development of Yellow-naped Amazon leading to F1 and F2 generation progenies.



STAGES OF BREEDING	DESCRIPTION
DIL is implementing the C	LOSE BREEDING AND PRODUCTION SYSTEM as well as RATEGY in various stages of breeding when applicable.
Segregated Conditioning	Birds with varying bloodlines are placed in a flight aviary where they can socialize and or bond with other birds (same species). This stage is the maturation period. Usually it will take three (3) -five (5) years to attain reproductive maturity.
Sexing	All unrelated, mature and with good quality birds are subjected to laparoscopy, to check maturity of reproductive organs such as the ovary/testes. This procedure is required to ensure proper pairing of sexually mature individuals.
Segregated Pairing	 Sexually compatible male and female will be set-up in either a Large Portable Cage or inside a Breeding Aviary. Newly paired breeder will be monitored and any aggression between the pair will be taken into consideration for possible break-up to avoid injury of either the male or the female.
Segregated Breeding	 Nest boxes with nesting materials are installed for egg-laying and special diet rich in Calcium will be provided. During breeding season a daily check-up of nest-boxes is required. However, noise and any other form of disturbance is avoided if not minimized.
Artificial Incubation	 Eggs are carefully pulled-out from nest boxes for cleaning and disinfection after which the eggs are placed inside a Grumbach Egg Incubator for artificial incubation Monitoring is done daily. Incubation is between 25-28 days
Hand-Rearing	Hatchlings are carefully cleaned and placed inside AICU-Animal Intensive Care Unit at the Nursery. A culture test of the eggshell is required if the hatclings exhibited any abnormal condition during incubation or immediately after hatching. Specially formulated diet will be given from day one until the bird is fully weaned. Weaning is between four (4)—eight (8) months.

10. IF THE OPERATION HAS ONLY BRED THE SPECIES TO THE FIRST GENERATION, DOCUMENTATION SHOWING THAT THE HUSBANDRY METHODS ARE THE SAME AS, OR SIMILAR TO THOSE THAT HAVE RESULTED IN SECOND-GENERATION OFFSPRING ELSEWHERE.

Not Applicable – The facility was able to produce up to second (F2).

11. PAST, CURRENT, AND EXPECTED ANNUAL PRODUCTION OF OFFSPRING, TOGETHER WITH THE INFORMATION ON THE PERCENTAGE OF: a) Female producing offspring each year b) Unusual fluctuations in the annual production of offspring (including an explanation of the probable cause).

Year	Total Percer	ntage of Produ	ctive Female		ction Acco Generation		Total Production
Parental Breeding Pair (12.12)	F1 Breeding Pair (16.16)	F2 Breeding Pair (N/A)*	F1	F2	F3		
1001	0	0	0	0	0	0	0
1981 1982	0	0	0	0	0	0	0
1982	0	0	0	0	0	0	0
1984	0	0	0	0	0	0	0
1985	0	0	0	0	0	0	0
1986	0	0	0	0	0	0	0
1987	0	0	0	0	0	0	0
1988	0	0	0	0	0	0	0
1989	8	Ö	0	2	0	0	2
1999	8	0	0	2	0	0	2
1990	17	0	0	4	0	0	4
1992	25	0	0	5	0	0	5
1992	17	0	0	3	0	0	3
1994	42	0	0	8	0	0	8
1995	50	0	0	10	0	0	10
1996	33	13	0	6	2	0	8
1997	25	0	0	3	0	0	3
1998	8	0	0	2	0	0	2
1999	0	6	0	0	1	0	1
2000	0	6	0	0	1	0	1
2001	0	50	0	0	17	0	17
2002	0	56	0	0	24	0	24
2003	0	36	0	0	34	0	34
2004	0	63	0	0	29	0	29
2005	0	75	0	0	33*	0	33
2006	0	75	0	0	37*	0	37

Legend = * indicates projected value

12. AN ASSESSMENT OF THE ANTICIPATED NEED FOR, AND SOURCE OF, ADDITIONAL SPECIMENS TO AUGMENT THE BREEDING STOCK TO INCREASE THE GENETIC POOL OF THE CAPTIVE POPULATION IN ORDER TO AVOID ANY DELETERIOUS INBREEDING.

Our current population of F1P-BR and the additional F2P-BR breeding stocks are sufficient sources of bloodlines to sustain our present captive breeding operation. Proper management of the progenies bloodlines through the use of a specimen control number using computerized relational database guides BII personnel to avoid pairing of related birds.

BII selects unrelated, mature and superior quality captive bred produced at the center. Contacts with other breeding operations worldwide allow the trade and exchange of this species thus eliminating the need to collect this species from the wild population.

13. TYPE OF PRODUCT EXPORTED

The products exported are of live captive-bred birds. Each bird is individually marked, either by closed leg-band or by microchip implant and inspected by the technical staff of the Philippine Protected Areas and Wildlife Bureau (PAWB) which keeps a record of each bird bred from the time the egg hatched.

14. DETAILED DECRIPTION OF THE MARKING METHODS USED FOR THE BREEDING STOCK AND OFFSPRING AND FOR THE TYPES OF SPECIMENS THAT WILL BE EXPORTED.


14.1 CLOSED LEG-BAND and OPEN LEG-BAND

A closed stainless steel leg-band with unique number series is used for all captive-bred progenies and breeding stocks at the center. Closed leg-band is usually attached ten to fifteen days (10-15) after hatching. The closed leg-band identification is being used by the Philippines CITES-CMA to identify the birds intended for export. Also this is the basis for the issuance of CITES export permit.

Close Leg-bands & Open Leg-band (1/2")








An open stainless steel leg-band is attached additionally to a female captive bred-progeny after surgical sexing. This will clearly identify the surgically sexed female because it is fitted with two leg-bands, a closed band on the right foot and an open band on the left.

14.2 MICROCHIP IMPLANT

An ISO microchip implant is also used for the identification of captive-bred progenies intended for export.



15. DESCRIPTION OF THE INSPECTION AND MONITORING PROCEDURES TO BE USED BY THE CITES MANAGEMENT AUTHORITY TO CONFIRM AND IDENTIFY THE BREEDING STOCK AND THE OFFSPRING AND TO DETECT THE PRESENCE OF UNAUTHORIZED SPECIMENS HELD AT OR INCORPORATED WITH THE OPERATION OR PROVIDED FOR EXPORT.

The following procedures are being adopted by the Protected Areas and Wildlife Bureau (PAWB), the CITES Management Authority in monitoring the captive breeding operations of BII;

- The collection of wildlife species from the natural habitat for breeding purposes is allowed only under a Wildlife Collector's Permit (WCP) issued by the PAWB. Exotic species maybe acquired from breeders duly accredited and registered with PAWB or through importation under an import permit issued by the said agency;
- 2. A Wildlife Farm Permit (WFP) is also required for the establishment of breeding farms in the country. A WFP holder is required to maintain and provide breeding facilities suitable for the avian species to be maintained and spacious enough to ensure the welfare of the birds. Likewise, leg bands for the identification of the breeders and progenies are being checked/verified by the CMA.
- 3. The breeding farm is required to maintain a record for each species maintained in the farm, which shall include the data specified below. The record is periodically inspected and the stocks verified by the CMA. Only the recorded/validated animal stocks in the farm as registered in the book are allowed for trade/disposition.
 - founding/breeder stocks- species, origin, age (date laid and date hatched) generation, sex, quantity, source, marking (leg-band, microchip etc) parental leg-band number if any.
 - b. progenies species, origin, generation, age (date laid and date hatched) sex, quantity, parental leg-band number.
 - additional acquisition local purchase, importation, exchange or donation, species, source, generation, origin,

- age, quantity, parental leg-band, receipt, permit if imported
- mortality /accidental escape species, source, generation, origin, age, generation, quantity, parental leg-band number
- hospital, nursery, incubator apparatus, and nesting area shall provide the same information as above.
- 4. The movement of the birds from the breeding farm to any point within the territorial jurisdiction of the Philippines or from the collection area to the farm should be accompanied by a local transport permit issued by the concerned field office of the CMA;
- 5. Only captive-bred wildlife species/sub-species produced in the breeding farms authorized under WFP are allowed for trade. Birds intended for export are inspected/validated by PAWB and verified against the production report regularly submitted by the establishment to the CMA. Export permits are issued only when the results of the inspection and verification conform with the requirements of the CMA;
- 6. The establishment is also required to observe cleanliness and sanitation in the maintenance of the breeding farm and facilities. This is to prevent possible contamination and /or spread of pests and/or diseases, which will affect the survival of either the captive-bred specimens or other wildlife species which maybe found in the surrounding area.
- 7. Cleanliness and Sanitation- the breeding operation shall observe cleanliness and sanitation in the maintenance of the breeding farm and facility. This is to prevent possible contamination and /or spread of pests and/or diseases, which may affect the survival of both captive-bred, and the wild population of animals and other wildlife species in the area.
- 16. DESCRIPTION OF THE FACILITIES TO HOUSE THE CURRENT AND EXPECTED CAPTIVE STOCK, INCLUDING SECURITY MEASURES TO PREVENT ESCAPES AND/OR THEFTS. DETAILED INFORMATION SHOULD BE PROVIDED ON THE NUMBER AND SIZE OF THE BREEDING AND REARING ENCLOSURES, EGG INCUBATION CAPACITY, FOOD PRODUCTION OR SUPPLY, AVIALABILITY OF VETERINARY SERVICES AND RECORD KEEPING.

The list of equipment and facilities presented below play a very vital role in the husbandry management and breeding of all Yellow-naped amazon at Birds International.

Facilities at Birds International

Description	Number of Units	Description	Description				
Hatchery-I & II	2	Incubation	86m ²				
Nursery 1-V	5	Hand-Rearing	1700m²				
Hanging Cages			1.				
Small	30	Weaning/Rearing	.92m³/cage				
Medium	120	Weaning/Rearing	2.63m³/cage				
Large	60	Weaning/Rearing	4.46m³ /cage				
Mini-Flight	30	Conditioning Newly Weaned Birds	400m²				
Conditioning Flight	30	Conditioning Newly Weaned Birds	1750m ²				
Breeding Aviary	27	Breeding	1500m ²				
Portable Breeding Cages							
Small	270	Breeding	3m³/ cage				
Medium	250	Breeding	6m³/ cage				
Large	100	Breeding	8m³/cage				
Food Preparation	2	Food Production for Breeders/Conditioning	180m²				
*********	3	Patient Treatment/Recovery	577m ²				
Hospital	1	Specimen culture	13.5m ²				
Laboratory Warehouse	2	Storage	480m ²				
	2	Storage	70m²				
Cold Storage-Warehouse Administration Building	- i	Record Keeping/Documentation	252m²				
	4	Accommodation	1,462m ²				
Caretaker's Quarter	1	Caretaker's Dining Area	25m²				
Dining Hall	2	Treatment /Isolation Area	730m²				
Quarantine Building Laundry Building	1	Laundry	60m²				

Equipment at Birds International

Description	Quantity	Purpose
Egg Incubators	30	Artificial Incubation
Egg Candlers	5	Egg Development Indication
AICU (Animal Intensive Care Units)	271	Hand-rearing
Walk-In Freezer	1	Food Storage
Endoscope	2	Endoscopy/Surgical Sexing
Photo-microscope	4	Bacteria Indentification
Mechanical Convection Incubator	I	Bacteria Culture
Surgitron	1	Surgery
Airconditioning Units	50	Temperature Control
Humidifier	2	Humidity Control
Microchip Reader	3	Microchip Number Identification
Computers (CPU+Monitor)	7	Record Keeping
Power Generators (90 KVA)	3	Stand-by Power Supply
Water Filtration (UV- Light)	1	Water Disinfection
Fork-Lift	1	Garbage Collection/Transport
Power Sprayer	5	Cleaning Cages
Mini-Van	1	Bird Export
Dump Truck	1	Materials/Equipment Transport
Reflotron	1	Blood Value Analyzer
Centrifuge	2	Blood Sample Preparation
Gas-Anesthesia Unit	2	Inhaled-Anesthesia
Digital Camera (mpeg/jpeg)	2	Picture/Documentation
Refrigerators	15	Food Storage


HATCHERY EQUIPMENT AND FACILITIES




Hatchery I

Hatchery II


NURSERY EQUIPMENT AND FACILITIES



CONDITIONING, BREEDING & PORTABLE BREEDING FACILITIES




FOOD PREPARATION EQUIPMENT AND COLD STORAGE FACILITIES




Chiller Warehouse


HOSPITAL AND LABORATORY EQUIPMENT AND FACILITIES



HOSPITAL AND LABORATORY EQUIPMENT AND FACILITIES



Mechanical Convection Incubator



Anesthesia machine

WATER TREATMENT AND FILTRATION/ LAUNDRY FACILITIES



Laundry room Washing/Drying machine Water Filtration/ UV Light Treatment & Storage

STAND-BY POWER GENERATOR SET AND TRANSPORT VEHICLE



CARETAKER'S QUARTERS AND DINING HALL




ADMINISTRATION FACILITIES



Prevention of the possible escape of birds is one of the many considerations in a cage design at BII. Almost all the cages used at the center can be serviced without the need to go inside to deliver the food to the birds. Food and water dishes are provided with through an especially designed hatches that can be pulled out or slide out during feeding. This design makes the escape of the birds inside the cage almost impossible.

Food Tray




Power-Pressure Spraying



Cleaning of the cage can usually be achieved by pressure hosing from the outside of the cage.

Nest-boxes are designed and situated in such a position that they also can be examined without the need to enter the cage.

Nest Box-Outside View



Nest Box- Inside View



On few occasions, when it is necessary to enter the cage to catch and examine more closely the bird, a special safety cage or cage trap is used to enter the main cage and prevent any possibility of escape.


Birds inside the breeding aviaries are housed in a completely enclosed building from which they cannot escape, even if they should come out of their cages. Breeding aviaries are secured through a screened double door.

Great care is taken whenever birds are being transferred to different facilities. These birds are always carried inside a transfer box with lock before going out of the aviary and the box is never opened unless it is fully secured inside the destination aviary.

Cage Trap



Aluminum Transport Box



The advent of technological breakthrough in the field of PC (Personal Computers) in the early 1980's enhanced and changed the old master list. Presently, the company has a relational database system that aids its staff in all aspects of avian management. This database was further enhanced with the use of C++, SQL and Visual Basic programming language. The database system at present can alert user not to pair related progenies as it does not accept records and data of related progenies intended to be paired.

17. DESCRIPTION OF THE STRATEGIES USED BY THE BREEDING OPERATION OR OTHER ACTIVITIES THAT CONTRIBUTE TO IMPROVING THE CONSERVATION STATUS OF WILD POPULATION(S) OF THE SPECIES.

The company is committed to support any effort of the Philippine government or any foreign government and non-government organizations to improve conservation status of the Yellow-naped amazon. The company's vast experience on the conservation of the species like Spix's macaw and the Philippine Eagle to name a few will likewise offer a great advantage to any conservation effort that will be initiated on this particular species.

18. ASSURANCE THAT THE OPERATION SHALL BE CARRIED OUT AT ALL STAGES IN A HUMANE (NON-CRUEL) MANNER.

The current success of BII in captive breeding not to mention other Appendix 1 species is a clear manifestation that indeed breeding operation at BII is carried out with utmost and humane manner. The company's commitment to follow the existing Animal Welfare Act of the Philippines and the Wild Life Resources Conservation and Protection Act is also an assurance that the company will protect, care and treat all species in a humane and non-cruel manner.

Annex A Yellow Naped Amazon

Species	Legband	Sex Age		Source	Evidence of Le	Comments	
Count	Number		100,000	Country (ISO-Code)	Import Permit	Date Issued	
	-			•			
1	601	F	UNK	SG	1980-06-26	P-153	A
2	692	F	UNK	SG	1980-06-26	P-153	Α
3	691	F	UNK	SG	1980-06-26	P-153	A
4	690	F	UNK	SG	1980-06-26	P-153	Α
5	685	F	UNK	SG	1980-06-26	P-153	Α
6	679	F	UNK	SG	1980-06-26	P-153	Α
7	678	F	UNK	SG	1980-06-26	P-153	Α
8	665	F	UNK	SG	1980-06-26	P-153	Α
9	650	F	UNK	SG	1980-06-26	P-153	Α
10	648	F	UNK	SG	1980-06-26	P-153	Α
11	646	F	UNK	SG	1980-06-26	P-153	Α
12	188	F	UNK	SG	1980-06-26	P-153	Α
13	653	M	UNK	SG	1980-06-26	P-153	Α
14	652	M	UNK	SG	1980-06-26	P-153	A
15	649	М	UNK	SG	1980-06-26	P-153	Α
16	296	M	UNK	SG	1980-06-26	P-153	A
17	200	М	UNK	SG	1980-06-26	P-153	Α
18	197	М	UNK	SG	1980-06-26	P-153	Α
19	195	M	UNK	SG	1980-06-26	P-153	Α
20	193	M	UNK	SG	1980-06-26	P-153	Α
21	193	М	UNK	SG	1980-06-26	P-153	Α
22	192	M	UNK	SG	1980-06-26	P-153	Α
23	191	M	UNK	SG	1980-06-26	P-153	A
24	189	М	UNK	SG	1980-06-26	P-153	Α
25	2864	F	UNK	SG	1980-06-26	P-153	DQ
26	2860	F	UNK	SG	1980-06-26	P-153	DQ
27	2859	F	UNK	SG	1980-06-26	P-153	DQ
28	2858	F	UNK	SG	1980-06-26	P-153	DQ
29	2851	F	UNK	SG	1980-06-26	P-153	DQ
30	2849	F	UNK	SG	1980-06-26	P-153	DQ
31	2848	М	UNK	SG	1980-06-26	P-153	DQ
32	2840	M	UNK	SG	1980-06-26	P-153	DQ
33	2834	М	UNK	SG	1980-06-26	P-153	DQ
34	2819	U	UNK	SG	1980-06-26	P-153	DQ
35	2818	U	UNK	SG	1980-06-26	P-153	DQ
36	2817	U	UNK	SG	1980-06-26	P-153	DQ

Annex A Yellow Naped Amazon

Species	Legband	Sex	Age	Source	Evidence of Leg	gal Acquisition	Comments
Count	Number		915 5 ,73	Country (ISO-Code)	Import Permit	Date Issued	
0.7			LINIX	SG	1980-06-26	P-153	DQ
	2814	U	UNK	30	1900-00-20	1-100	UG
37	2814	U	UNK	SG	1980-06-26	P-153	DQ
38	2814 2812 2713	U	577000000				740,000

Legend

DQ =

Died during Quarantine

A =

Alive

SG=

Singapore

UNK =

Unknown

U=

Unsexed

Annex B Yellow Naped Amazon

Species	Legband	aband Sex		Source	Evidence of Leg	gal Acquisition	Comments
Count Number	Age	Country (ISO-Code)	Import Permit	Date Issued			
1	601	F	UNK	SG	1980-06-26	P-153	Α
2	692	F	UNK	SG	1980-06-26	P-153	Α
3	691	F	UNK	SG	1980-06-26	P-153	Α
4	690	F	UNK	SG	1980-06-26	P-153	Α
5	685	F	UNK	SG	1980-06-26	P-153	Α
6	679	F	UNK	SG	1980-06-26	P-153	Α
7	678	F	UNK	SG	1980-06-26	P-153	Α
8	665	F	UNK	SG	1980-06-26	P-153	Α
9	650	F	UNK	SG	1980-06-26	P-153	Α
10	648	F	UNK	SG	1980-06-26	P-153	Α
11	646	F	UNK	SG	1980-06-26	P-153	Α
12	188	F	UNK	SG	1980-06-26	P-153	Α
13	653	M	UNK	SG	1980-06-26	P-153	Α
14	652	M	UNK	SG	1980-06-26	P-153	A
15	649	М	UNK	SG	1980-06-26	P-153	Α
16	296	М	UNK	SG	1980-06-26	P-153	Α
17	200	М	UNK	SG	1980-06-26	P-153	Α
18	197	M	UNK	SG	1980-06-26	P-153	Α
19	195	М	UNK	SG	1980-06-26	P-153	Α
20	193	М	UNK	SG	1980-06-26	P-153	Α
21	193	M	UNK	SG	1980-06-26	P-153	A
22	192	M	UNK	SG	1980-06-26	P-153	Α
23	191	М	UNK	SG	1980-06-26	P-153	Α
24	189	M	UNK	SG	1980-06-26	P-153	Α

Legend

A =

Alive

SG =

Singapore

UNK =

Unknown

Annex C - Yellow Naped Amazon (Information on the Percentage of Mortalities in the Different Age Groups) Male / Female

F1= 45 heads. Total Production 1989-98

Mortalities	=	13 hds	29	%
Live	=	32 hds	71	%
Total	=	45 hds	100	%

Classification of Mortalities		Se	Total					
(Age Groups)	M		F		U			
(Age Groups)	Hds	%	Hds	%	Hds	%	Hds	%
	1 0	0		0	10	22	10	22
1 day 17 months				O	10	66	10	22
1day-12 months 12 months - 48 months	1	2	1	2	0	0	2	4

13 29

F2= 108 heads. Total Production 1996-Present 11.

Г	Mortalitie	s =	1 hd	1	%
	Live	=	107 hds	99	%
	Total	=	108 hds	100	%

	Se	ex Class	sificat	ion		To	Total	
M		F	F		J			
Hds	%	Hds	%	Hds	%	Hds	%	
0	0	0	0	1	1	1	1	
0	0	0	0	0	0	0	0	
0	0	0	0	0	0	0	0	
	Hds 0	M Hds %	M F Hds % Hds 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	M F Hds % Hds %	Hds % Hds % Hds 0 0 0 0 1 0 0 0 0 0	M F U Hds % Hds % 0 0 0 1 1 0 0 0 0 0 0	M F U Hds % Hds % Hds 0 0 0 1 1 1 0 0 0 0 0 0	

1980-06-26

Metro-Manila Date:

EMPORT PERMIT NO. P-153

Mr. Antonio de Dios 99 Timog Ave.

Sir/Madam:

Pursuant to your request of even date 4 you are nereby given permission to impact forty (40) parrots From Singapore that the Philippines for breeding purposes, subject to the proceeding of paragraphs 22 and 24 of Animal Industry Administrative Order No. 7-2, quoted hereunder:

IMPORTATION OF FOWLS: - Fowls shall be accompanied by a health contificate issued by the proper veterinary authority at the port of emparkation, certifying to the fact:

GERTIFIED TRUE COPY

- That the fowle are free from and have not been in contact with or expected as any dangerous and communicable disease arTwoting have at least sixty (60) days before the data of shipment.
- That the finds have been tested against pullorum ten (10) days before objement with negative results. In case of stacks a certification of flock from whom they come from for freedom from pullorum will cuffice.
- Phas the found have come from flocks that are from from pullorum and 'ankosis complex.
- 24. DISPOSAL OF INFLUENCE FOWES: Upon arrival of the said flowls at the port of entry, they shall be placed under quarantine at any place dorighated by the Director of Animal Industry and subject to sall orum test by the representative of this Burrelli.

Aspould and Fig. of fouls arrive infected with any dangerous communicable bisease, all of the sick fouls shall be destroyed or LShould ah cremated and those appearant, well ones be held under quarantine. He compensation Shall be allowed for the chickens destroyed.

O. R. No. The permittee whill pay the the Boreau of Animal Endustry the tol-

- For the essign of this import Permits / 2.00 1st 20 hds. _____ Sec. VII, Animal Industry Adm. order No. 6-18
- b) For the inspecutor and issuance of landing permit upon arrivel of the said birds at the port of entry-7 5.00 1st 5 hds. & .50 each for excess I, An. Ind. Adm. Order No. 6-14)

Jul. 30, 1980 This permit stall expure on and is subject to cancellation should any dangerous communicable poultry disease break out at the place of uriols, or may be revoked at anytime before said date if the interest of the pavernment so requires.

Very truly yours,

DALMACIOTA CEPTED THE KARLESTEE HARD IN

rhe/7-17-79 Dr. Pastor Cruz-OIC, MIA

